

CERTIFICATE OF ANALYSIS

EcoRI

#ER0275 10000 u

Lot: Expiry Date:

5'...G↓A A T T C...3'
3'...C T T A A↑G...5'

Concentration: 10 u/μl
Source: *E.coli* that carries the cloned *ecoRI* gene from *Escherichia coli* RY13
Supplied with: 4x1 ml of 10X Buffer EcoRI
1 ml of 10x Buffer Tango™

Store at -20°C



In total 6 vials.

BSA included: Lot# BSA62-313P

RECOMMENDATIONS

1X Buffer EcoRI (for 100% EcoRI digestion)

50 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 100 mM NaCl, 0.02% Triton X-100, 0.1 mg/ml BSA.

Incubation temperature

37°C.

Unit Definition

One unit is defined as the amount of EcoRI required to digest 1 μg of lambda DNA in 1 hour at 37°C in 50 μl of recommended reaction buffer.

Dilution

Dilute with the Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA and 50% glycerol.

Double Digests

Tango™ Buffer is provided to simplify buffer selection for double digests. 98% of Fermentas restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to the Fermentas Catalog or go to www.fermentas.com/doubledigest to choose the best buffer for your experiments.

1X Tango™ Buffer:

33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA.

Storage Buffer

EcoRI is supplied in: 10 mM potassium phosphate (pH 7.4 at 25°C), 300 mM NaCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA, 0.15% Triton X-100 and 50% glycerol.

Recommended Protocol for Digestion

- Add:

nuclease-free water	16 µl
10X Buffer EcoRI	2 µl
DNA (0.5-1 µg/µl)	1 µl
EcoRI	0.5-2 µl ^{a, b)}
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours^{b)}.

The digestion reaction may be scaled either up or down.

Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 µl (~0.1-0.5 µg of DNA)
nuclease-free water	18 µl
10X Buffer EcoRI	2 µl
EcoRI	1-2 µl ^{a, b)}
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours^{b)}.

^{a)} This volume of the enzyme is recommended for preparations of standard concentrations (10 u/µl), whereas HC enzymes (50 u/µl) should be diluted with the Dilution Buffer to obtain 10 u/µl concentration.

^{b)} See Note.

Thermal Inactivation

EcoRI is inactivated by incubation at 65°C for 20 min.

ENZYME PROPERTIES

Enzyme Activity in Fermentas REase Buffers, %

EcoRI	B	G	O	R	Tango™	2X Tango™
100	0-20	NR	100	100*	NR	100

*Star activity appears at a greater than 5-fold overdigestion (5 u x 1h).

NR: buffer is not recommended, because of high star activity.

Methylation Effects on Digestion

Dam: never overlaps – no effect.

Dcm: never overlaps – no effect.

CpG: may overlap – cleavage impaired.

EcoKI: never overlaps – no effect.

EcoBI: may overlap – no effect.

Stability during Prolonged Incubation

A minimum of 0.2 units of the enzyme is required for complete digestion of 1 µg of lambda DNA in 16 hours at 37°C.

Digestion of Agarose-embedded DNA

A minimum of 5 units of the enzyme is required for complete digestion of 1 µg of agarose-embedded lambda DNA in 16 hours.

Compatible Ends

XapI, MunI, TasI

Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
5	0	1	1	1	1	1

Note

A large excess of EcoRI (7.5 u/µg DNA x 16 hours), low salt concentration, high pH, or the replacement of Mg²⁺ by Mn²⁺ may result in star activity.

For **QUALITY CONTROL ASSAY DATA** see back page

QUALITY CONTROL ASSAY DATA

Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 80-fold overdigestion with EcoRI (5 u/μg lambda DNA x 16 hours).

Ligation/Recutting Assay

After a 50-fold overdigestion (3 u/μg DNA x 17 hours) with EcoRI, more than 95% of the digested DNA fragments can be ligated at a 5'-termini concentration of 0.02 μM. More than 95% of these sites can be recut.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of EcoRI for 4 hours.

Blue/White Cloning Assay

pUC57 was incubated with 10 units of EcoRI for 16 hours. After religation and transformation, the background level of white colonies was 0.1%.

Quality authorized by:



Jurgita Zilinskiene

PRODUCT USE LIMITATION.

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.fermentas.com for Material Safety Data Sheet of the product.